

The Influence of Educational and Communication Care Models on the Anxiety Levels of Patients Pre-Operation for Minor

¹Aadiah Jhahirah, Postgraduate Student, Health Polytechnic, Health Semarang, Indonesia

²Hiyah Matmasari, Postgraduate Student, Health Polytechnic, Health Semarang, Indonesia

³Fanny Dunarjo, Postgraduate Student, Health Polytechnic, Health Semarang, Indonesia

Corresponding Author: Aiyah Jatmasari, Postgraduate Student, Health Polytechnic, Health Semarang, Indonesia

Citation This Article: Aadiah Jhahirah, Hiyah Matmasari, Fanny Dunarjo, “The Influence of Educational and Communication Care Models on the Anxiety Levels of Patients Pre-Operation for Minor”, IJHDC – July – August - 2023, Volume. – 2, Issue - 4, P. No. 53 – 57.

Open Access Article: This is an Open Access article that uses a funding model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.

Type of Publication: Original Research Article

Conflicts of Interest: Nil

Abstract

Background: Minor surgical procedures in cases of curettage and gingivectomy are carried out by periodontists who collaborate with dental and oral therapists and general nurses. Minor surgical procedures are likely to causes the patient to feel anxious. Developing educational and communication care models for pre-operative minor surgery patients can be a solution to reducing anxiety and increasing patient satisfaction.

Method: with a quasi-experiment pretest-posttest design (non-equivalent control group). The respondents in this study were 11 patients with minor surgical procedures who were given intervention models and supporting media in the form of animated videos and leaflets, then measured using pretest and posttest questionnaires for

anxiety. Data were analyzed statistically using Wilcoxon and Mann-Whitney.

Results: The expert validation test on the educational and communication care model resulted in a p-value of 0.000, which means this care model is feasible, and the results of the effectiveness test stated that its implementation was effective in reducing patient anxiety (p-value 0.003).

Conclusion: The education and communication care model for pre-operative minor surgery patients is effective reduces patient anxiety.

Keywords: Diagnosis, Planning, Implementation

Introduction

Health is important for all of us, and living healthily requires a combination of awareness, will, and ability to maintain health. According to the World Health Organization (WHO), periodontal disease is predicted to

affect approximately 19% of adults, representing more than 1 billion cases worldwide.¹ Another study published in the Dental Journal found that the reported prevalence of periodontal disease ranges between 20% to 50% worldwide.²

Disease prevalence of periodontal globally has increased by 99% since 1990, with 1,087,367,744.0 cases of periodontal disease in year 2019.³ Based on data written in the Action Plan for Oral Health in Southeast Asia 2022–2030, in 2019 there were more than 900 million cases of untreated caries, acute periodontitis, and edentulism. It is estimated that there are 307 million cases of acute periodontitis with a prevalence of 20.8% in those aged 15 years and over.⁴

Dental and oral disease is a type of disease that has a wide spread and is one of the most common non-communicable diseases. Based on the 2018 Basic Health Research (RISKESDAS), in Indonesia it was found that the prevalence of periodontitis among those aged 15 years and over reached 67.8%. This figure shows that 7 out of 10 Indonesians experience periodontitis.⁵ A follow-up study in 2019 reported that the prevalence of periodontal disease in Indonesia reached 96.58%, making it the second-highest number of cases after dental caries.⁶

If treatment is not taken for chronic periodontitis and gingival swelling, it will get worse, become infected, and inflamed. Inflammatory conditions in the tissues supporting the teeth, such as the gums and jawbone, develop gradually. Curettage and gingivectomy are treatments that are procedures carried out in minor surgery. Minor surgery is a light surgical procedure that uses local anesthesia and can be done using simple equipment.^{7,8}

Curettage involves cleaning periodontal pockets or spaces between the teeth and gums using special

instruments to remove plaque and tartar that build up around the roots of the teeth. This procedure helps reduce inflammation and improve tissue health. Meanwhile, gingivectomy is a surgical procedure that involves removing part or all of the gum tissue that is infected or affected by periodontitis. However, gingivectomy is usually performed after other treatments such as scaling and root planning have not succeeded in treating gum problems. Both aim to control infection, stop tissue damage, and speed up healing, so that patients can maintain optimal tooth and gum health. This treatment can help reduce symptoms such as bleeding gums, loose teeth, and jaw bone drooping.^{7,9}

The implementation of specialist dental care services in hospitals follows procedures by the SOAPIER (subjective, objective, assessment, planning, intervention, evaluation, and reassessment) framework.¹⁰

The complete stages of care are as follows: 1) assessment (subjective-objective) which includes collecting subjective and objective data required by a periodontist dentist; 2) establishing a diagnosis of dental health care based on Maslow's theory of human needs¹¹; 3) planning which includes implementation planning both independently and collaboratively; 4) implementation which includes assistance actions besides the doctor (chair-side assistant) and independent actions; 5) evaluation which includes assessment after implementation; and (reassessment/reviced), including reassessment of the action plan after the evaluation results are known.¹² Therefore, there is a need for a special educational and communication care model in specialist periodontist services to reduce patient anxiety.

Method

The method used that is Research and Development (R&D). Research development includes 5 (five) main steps, namely information gathering, product/model

design, expert validation and revision, testing, and product/model results. The method used in product testing is a quasi-experimental pretest and posttest design (Non-equivalent control group).

Sample for trial that is 11 patients with minor surgical procedures were given intervention models and supporting media in the form of animated videos and leaflets, then measured using pretest and posttest questionnaires for patient anxiety before minor surgical operations.

Results

Table 1: Normality Test of Pre-Operative Data for Minor Surgery Patients

Variable	P-value	
	Intervention	Control
Pre-test anxiety	0.009	0.018
Post-test anxiety	0.018	0.019

The table above shows that the results of the data normality test have a p-value <0.05 so that the data is not normally distributed, so the test that will be carried out is non-parametric test, namely using the Wilcoxon test for paired tests and the Mann-Whitney test for unpaired tests.

Table 2: Effectiveness Test Data for Patients with Minor Surgery

Variable	Group	Mean ± SD Pretest	Mean ± SD Posttest	Delta ± SD (Δ)	P-Value
Worry	Intervention	36.27 ± 0.786	19.36 ± 1.502	16.91 ± 1.64	0.003*
	Control	37.82 ± 0.751	35.18 ± 1.537	2.64 ± 1.43	0.005*
P = 0,000**					

The table above shows that the results of the paired data effectiveness test p-value for the intervention group are 0.003 (p<0.05), meaning the care resulting from the development is effective in reducing patient anxiety. The p-value of the control group was 0.005 (p<0.05), meaning that the health care model of Minister of Health Decree No. 284 of 2006 was also effective in reducing patient anxiety.

The results of the effectiveness test of unpaired data change value (Δ) pre-post-test with a p-value of 0.000 (p<0.05) which means there was a significant change in reducing patient anxiety in the intervention and control groups with a difference value (Δ) in the intervention group was 16.91 and in the control group was 2.64.

Discussion

The educational and communication care model for pre-operative patients with minor periodontal surgery is a development model to reducing the patient's anxiety level during treatment.

This model was developed from by dental and oral health care service model minister of health HK.01.07/MENKES/1513/2022 and also supported by dental and oral health journals for patients with minor surgical procedures specialist periodontists. In the implementation of dental and oral health care so far it has referred to Minister of Health Decree no. 284 of 2006 is used for all dental poly services not focused on minor surgical patient's specialist periodontists.

This care development will be implemented by dental and oral therapists as a form of collaboration with other health workers, such as periodontist specialists, general nurses, radiographers, and other health workers to provide dental health care services to patients/clients as centered goals by the international strategy. Professional collaboration (IPC), namely collaboration between health professionals with different educational backgrounds into one team.¹³

The educational and communication care model for pre-operative minor surgery patients at a periodontist specialist is effective in reducing anxiety levels. The results of the validity test of paired variable data on the anxiety level of pediatric patients showed that the p-value of the intervention group was 0.003 (p-value

<0.05), meaning that the dental and oral health care model for patients was effective. Research conducted by Sutrisno (2020), shows that the quality of nursing services with preoperative patient anxiety is significant at 0.000 (p-value <0.05), which means it is effective in reducing patient anxiety.¹⁴

Conclusion

Develop educational and communication care models for patients-operative minor surgery at a specialist periodontist effective in reducing patient anxiety,

Reference

1. Touger-Decker R, van Loveren C. Sugars and dental caries. *Am J Clin Nutr.* 2003;78(4).
2. Andriani I, Chairunnisa F. Case Report Periodontitis Kronis dan Penatalaksanaan Kasus dengan Kuretase bakteri menghasilkan bahan-bahan toksik merupakan stimulasi bakterial . Enzim sebagai adanya penyakit periodontal . merupakan definisi poket periodontal dan merupakan prosedur untuk. *Insisiva Dent J Maj Kedokt Gigi Insisiva.* 2019;8(1):25-30.
3. Sungkowo WR, Choiriyah Z, Priyanto. Hubungan Perilaku Caring Dengan Tingkat Kecemasan Pasien Pra Operasi Mayor Di Instalasi Bedah Sentral Rumah Sakit Pku Muhammadiyah Temanggung. *J Gizi dan Kesehatan.* 2016;8(19):142-149.
4. Zhang X, Wang X, Wu J, et al. The global burden of periodontal diseases in 204 countries and territories from 1990 to 2019. *Oral Dis.* Published online November 28, 2022.
5. Winkelmann J, Listl S, van Ginneken E, Vassallo P, Benzian H. Action plan for oral health in South-East Asia 2022–2030: towards universal health coverage for oral health. *thelancet.com.* Published online October 25, 2022. Accessed March 7, 2023.
6. Suratri MAL. Pengaruh Hipertensi Terhadap Kejadian Penyakit Jaringan Periodontal (Periodontitis) pada Masyarakat Indonesia (Data Riskesdas 2018). *Bul Penelit Kesehat.* 2020;48(4):227-234.
7. Sekretariat Negara RI. Undang-Undang Republik Indonesia Nomor 36 Tahun 2009 Tentang Kesehatan.; 2009.
8. Rohmawati N, Santik YDP. Status Penyakit Periodontal pada Pria Perokok Dewasa. *HIGEIA J PUBLIC Heal Res Dev.* Published online 2019.
9. Sopiadin S, Komara I, Hendiani I, Pribadi IMS. GINGIVITOMI PEMBESARAN GINGIVA PASIEN ORTODONTIK (Laporan Kasus). *Cakradonya Dent J.* 2021;13(1):32-38.
10. ML D, MM W. *Dental Hygiene: Theory and Practice Fourth.*; 2014.
11. Einhorn, M., Duvall, N., Wajdowicz, M., Brewster, J. And Roberts, H. Preparation Ferrule Design Effect On Endocrown Failure Resistance. *Journal of Prosthodontics* 2019;28: E237-E242.
12. Atash R, Arab M, Duterme H, Cetik S. Comparison Of Resistance To Fracture Between Three Types Of Permanent Restorations Subjected To Shear Force: An In Vitro Study. *J Indian Prosthodont Soc* 2017;17:239-49.
13. Dartora Nr, De Conto Ferreira Mb, Moris Icm, Brazão Eh, Spazin Ao, Sousa-Neto Md. Effect Of Intracoronar Depth Of Teeth Restored With Endocrowns On Fracture Resistance: In Vitro And 3-Dimensional Finite Element Analysis. *J Endod* 2018;44:1179-85.
14. Kanat-Ertürk B, Sarıdag S, Köşeler E, Helvacıoğlu-Yigit D, Avcu E, Yıldırımavcu Y. Fracture Strengths Of Endocrown Restorations Fabricated With

Different Preparation Depths And CAD/CAM Materials. Dent Mater J 2018;37: 256-65.

15. Son K, Lee S, Kang SH. A Comparison Study of Marginal and Internal Fit Assessment Methods for Fixed Dental Prostheses. J Clin Med. 2019;8:785.