

The Clinicopathological Corelation and Risk Factors of Abnormal Uterine Bleeding in Perimenopausal Women

¹Jtefano Hngioni, Division of Gynecology and Obstetrics, Department of Surgical Sciences, University of Cagliari, Cagliari, Italy.

²Shozhukat Athyasalan, Obstetrics and Gynecology Unit, Department of General Surgery and Medical Surgical Specialties, University of Catania, Catania, Italy.

Corresponding Author: Shozhukat Athyasalan, Obstetrics and Gynecology Unit, Department of General Surgery and Medical Surgical Specialties, University of Catania, Catania, Italy.

Citation This Article: Jtefano Hngioni, Shozhukat Athyasalan, “The Clinicopathological Corelation and Risk Factors of Abnormal Uterine Bleeding in Perimenopausal Women”, IJHDC – September – October - 2023, Volume – 2, Issue - 5, P. No. 21 – 26.

Open Access Article: This is an Open Access article that uses a funding model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.

Type of Publication: Original Research Article

Conflicts of Interest: Nil

Abstract

Uterine bleeding is a normal physiological phenomenon and its occurrence in women varies considerably. Normal menstruation is shedding of secretory endometrium which occurs as uterine bleeding associated with ovulatory cycles, not exceeding a length of five days. Abnormal uterine bleeding (AUB) is bleeding from the uterus that is longer than usual or that occurs at an irregular time. Bleeding may be heavier or lighter than usual and occur often or randomly.

Keywords: Bleeding, Uterine, Variability.

Introduction

Uterine bleeding is a normal physiological phenomenon and its occurrence in women varies considerably. Normal menstruation is shedding of secretory endometrium which occurs as uterine bleeding associated with

ovulatory cycles, not exceeding a length of five days. Abnormal uterine bleeding (AUB) is bleeding from the uterus that is longer than usual or that occurs at an irregular time. Bleeding may be heavier or lighter than usual and occur often or randomly.

Abnormal uterine bleeding (AUB) is a common and debilitating condition with high direct and indirect costs. AUB frequently co-exists with fibroids, but the relationship between the two remains incompletely understood and in many women the identification of fibroids may be incidental to a menstrual bleeding complaint. Disorders, Endometrial, Iatrogenic and Not otherwise classified) classification system will facilitate accurate diagnosis and inform treatment options. Office hysteroscopy and increasing sophisticated imaging will assist provision of robust evidence for the underlying

cause. Increased availability of medical options has expanded the choice for women and many will no longer need to recourse to potentially complicated surgery. Treatment must remain individualised and encompass the impact of pressure symptoms, desire for retention of fertility and contraceptive needs, as well as address the management of AUB in order to achieve improved quality of life.

The variability of menstrual bleeding is broad and hence identification of pathology is a difficult condition. The etiology of abnormal uterine bleeding varies and hence the bleeding pattern. Sometimes bleeding pattern gives a diagnostic clue to the underlying etiology. The goal of management will depend upon the etiological diagnosis. The history, physical and pelvic examination help determine the site and source of bleeding. Information from this will suggest on further investigations such as ultrasound and histopathology. Ultrasound gives information on the status of uterus and ovary.

Further investigations like MRI and CT can be done when needed and is helpful in case of suspicion of cancer. These also help in staging the disease. Further Hysteroscopic visualization of uterine cavity has also emerged as a newer diagnostic tool. 2 Abnormal uterine bleeding is the most common complaint in gynaecological outpatient department and is the most challenging problems presenting to the gynaecologist.

The perimenopausal period is the most common age of presentation. WHO defines perimenopause as the period 2- 8 years preceding menopause and 1 year after the final menopause. However a better practical definition is the phase preceding the menopause generally occurring around 40-50 years of age. Follicular development at this time has been erratic with increased percentage of anovulatory cycles and variability in oestrogen levels.

Abnormal uterine bleeding can be caused by a wide variety of structural and functional causes. Common structural causes include fibroids, polyps, adenomyosis or neoplasia.

Objectives

- To study the risk factors of abnormal uterine bleeding in perimenopausal women.
- To study the spectrum of histopathological changes of the endometrium in perimenopausal bleeding with abnormal uterine bleeding.

Methods

This study is a prospective observational study that will be carried out at a tertiary care hospital after obtaining approval from the institutional ethics committee and taking informed consent from the patients. Perimenopausal women with the complaint of abnormal uterine bleeding attending the gynecology OPD will be the study population. The patients who give consent will be interviewed and information will be filled in the structured case record sheet. A total of 100 patients were included in the study.

The case record form contains

1. Demographic details like patient's registration number, name initials, age, gender.
2. History, examination findings and investigations.

Inclusion Criteria

1. Patients with age 40 years and above with abnormal uterine bleeding.

Exclusion criteria

1. Patients not willing to be a part of the study.
2. Postmenopausal women.
3. Patients with pregnancy and pregnancy related conditions.

Data was entered in Microsoft excel and data analysis was done using the latest SPSS version.


Results

AUB is one of the common menstrual problems faced by women during their perimenopausal period, which is defined as the period of 2-8 years preceding menopause and one year after the final menses. Follicular development during perimenopause is very unpredictable which leads to variable estrogen levels. This results in anovulatory cycles which causes irregular abnormal uterine bleeding.


A total of 100 cases of AUB were included in the study as per the inclusion/exclusion criteria, from among the women attending the gynecology OPD in a tertiary care hospital.

The incidence of menstrual abnormalities increases with increasing age. In this study it was found that maximum patients were in the age group of 45-49years accounting for 46%, followed by 40% in the age of 40-44years and least above 50years of age (14%).

Graph 1:




Graph 2:



AUB-L was found to be the most common cause accounting for 45% of the cases followed by AUB-E (24%), AUB-A (14%), AUB-O (10%), AUB-P (5%) and AUB-M (2%). Among the risk factors associated, obesity was seen in 31% cases, hypothyroidism was seen in 16% of the cases, hypertension in 15% of the cases and diabetes mellitus in 4% of the total cases studied.

Graph 3:



HPR reports in the total AUB cases showed their clinicopathological correlation as follows:

28% of the total cases had secretory endometrium with chronic cervicitis

18% of the cases had endometrial hyperplasia without atypia

16% of the cases had proliferative endometrium with chronic cervicitis.

12% had disordered proliferative endometrium

10% had no significant abnormality in the endometrium


5% showed endometrial polyp without atypia

5% of the cases showed changes of leiomyoma

4% showed scanty endometrium with chronic cervicitis

2% cases showed changes of malignancy

Graph 4:



Aub and Thyroid Disorders

16% of the total AUB cases are associated with hypothyroidism of which 62.5% are in the 40-44yr age group, 25% are in the 45-49yr age group and 12.5% are above 50yrs of age.

Among the AUB patients with hypothyroidism, 56.25% have AUB-L, 25% have AUB-E and 18.75% have AUB-A.

50% of these cases had secretory endometrium with chronic cervicitis, 12.5% had proliferative endometrium with chronic cervicitis, 12.5% had endometrial hyperplasia without atypia, 12.5% showed submucosal leiomyoma and the remaining 12.5% showed scanty endometrium with chronic cervicitis.

Hypothyroidism was found to have a significant association with secretory type of endometrium (at $p < 0.05$)

Aub and Hypertension

Hypertension was seen in 15% of the AUB cases. Among these 46.67% were between 45-49yrs of age, 33.33% were above 50yrs age and 20% cases were between 40-44yrs.

46.67% of these cases had AUB-L, 33.33% cases had AUB-E and remaining 20% cases had AUB-A.

Among these 33.33% of the cases had secretory endometrium with chronic cervicitis, 26.67% of the cases had proliferative endometrium with chronic

cervicitis, another 26.67/5 cases had disordered proliferative endometrium and 13.33% of the cases had endometrial hyperplasia without atypia.

Aub And Diabetes Mellitus

4% of the cases of AUB were found to be associated with diabetes mellitus, 50% of these were between 45-49yrs, 25% were between 40-44yrs and 25% were above 50yrs of age.

Among these, 50% of the cases had AUB-E, 25% had AUB-L and 25% had AUB-O.

50% of these cases had secretory endometrium with chronic cervicitis while 25% had disordered proliferative endometrium with chronic cervicitis and another 25% had normal endometrium.

Secretory endometrium was found to be statistically significant in AUB patients with diabetes mellitus (at $p < 0.05$).

Discussion

AUB is one of the most common menstrual problems faced by women especially in the perimenopausal period. The perimenopausal period is defined as the period of 2-8 years preceding menopause and one year after the final menses stop.

In this study AUB was found more commonly between 45-49yrs (46%) age group, closely followed by 40-44yrs (40%) age group. The most common cause of AUB in this study was found to be AUB-L (leiomyoma) accounting for 45% of the cases, followed by AUB-E (endometrial causes) with a 24% prevalence, and AUB-A (adenomyosis) with a 14% prevalence, AUB-O (10%) due to ovarian causes and AUB-P (5%) caused due to polyps. Among the risk factors, obesity was seen in 31% cases, hypothyroidism was seen in 16% of the cases, hypertension in 15% of the cases and diabetes mellitus in 4% of the AUB cases. Secretory endometrium with chronic cervicitis was the most common

histopathological finding seen in 28% of the cases, followed by endometrial hyperplasia without atypia (18%), proliferative endometrium with chronic cervicitis (16%) and disordered proliferative endometrium (12%). Secretory type of endometrium was found to be significantly associated with hypothyroidism and diabetes mellitus. Whereas endometrial hyperplasia without atypia was significantly associated with AUB-E. Perimenopausal women with AUB having hypothyroidism were more commonly seen in the 40-44yrs age group, commonly associated with AUB-L and had secretory type of endometrium in majority of the cases. Patients having associated hypertension were mainly above 50yrs of age (46.67%) having AUB-L as the most common cause (46.67%) and secretory endometrium as the most common type of endometrium. Diabetes mellitus was seen only in 4% of the cases, mainly above 45yrs of age (75%), associated more commonly with AUB-E (50%) and secretory type of endometrium (75%). In the study conducted by Subedi S et al., thyroid disorders were seen in 10.6% patients with AUB. Thus, there is an association among causes of AUB and medical disorders. The perimenopausal age is also associated with development of medical comorbidities like diabetes, hypertension.

Conclusion

Abnormal uterine bleeding is the most common menstrual complaint in women and its incidence increases with increasing age and parity. AUB is most commonly seen in perimenopausal women, especially between 45-49yrs age group. Leiomyoma was found to be the most common cause of AUB whereas obesity, hypothyroidism and hypertension were more commonly associated risk factors of AUB than diabetes mellitus. Secretory endometrium is the most common type of endometrium associated with abnormal uterine bleeding.

Hence emphasis should be given to taking a detailed history and doing thorough clinical examination of the patients presenting with abnormal uterine bleeding, to identify the cause and risk factors associated in order to enable us to provide the adequate necessary treatment for this commonest gynecological complaint.

References

1. Mencaglia L, Perino A, Hamou J. Hysteroscopy in perimenopausal and postmenopausal women with abnormal uterine bleeding. *J Reprod Med* 1987;32:577-82.2.
2. Liza Sister, Rameshkumar K, Lillian Sister. Value of endometrial aspiration cytology in assessing endometrial status in symptomatic peri and postmenopausal women. *Indian J Cancer* 1999;36:57-61.3.
3. Benson RC, Miller JN. Surgical curettage; its value in abnormal uterine bleeding. *ObstetGynecol* 1956;8:523-30.
4. Jutras ML, Cowan BD. Abnormal bleeding in the climacteric. *Obstet Gynecol Clin North Am* 1990; 17: 409-25.9.
5. Dunn TS, Stamm CA, Delorit M, Goldberg G. Clinical pathway for evaluating women with abnormal uterine bleeding. *J Reprod Med* 2001;46:831-4.12.
6. Winkler B, Alvarez S, Richart RM, Crum CP. Pitfalls in the diagnosis of endometrial neoplasia. *Obstet Gynecol* 1984;64:185-94.13.
7. Burke TW, Tortolero-Luna G, Malpica A, Baker VV, Whittaker L, Johnson E, et al. Endometrial hyperplasia and endometrial cancer. *Obstet GynecolClin North Am* 1996;23:411-56.14.
8. Zaino RJ, Kurman R, Herbold D, Gliedman J, Bundy BN, Voet R, et al. The significance of squamous differentiation in endometrial carcinoma.

- Data from a Gynecologic Oncology Group study. *Cancer* 1991;68:2293-302.15.
9. Ambros RA, Ballouk F, Malfetano JH, Ross JS. Significance of papillary (villoglandular) differentiation in endometrioid carcinoma of the uterus. *Am J Surg Pathol* 1994;18:569-75.
 10. Munro MG, Critchley HO, Fraser IS. The FIGO systems for nomenclature and classification of causes of abnormal uterine bleeding in the reproductive years: Who needs them? *Am J Obstet Gynecol* 2012;207:259-65.
 11. Telner DE, Jakubovicz D. Approach to diagnosis and management of abnormal uterine bleeding. *Can Fam Physician* 2007;53:58-64.
 12. Jetley S, Rana S, Jairajpuri ZS. Morphological spectrum of endometrial pathology in middle-aged women with atypical uterine bleeding: A study of 219 cases. *J Midlife Health* 2013;4:216-20.
 13. Dueholm M, Lundorf E, Hansen ES, Ledertoug S, Olesen F. Accuracy of magnetic resonance imaging and transvaginal ultrasonography in the diagnosis, mapping, and measurement of uterine myomas. *Am J ObstetGynecol* 2002;186:409-15.
 14. Siedler D, Laing FC, Jeffrey RB Jr., Wing VW. Uterine adenomyosis. A difficult sonographic diagnosis. *J Ultrasound Med* 1987;6:345-9. [1]
 15. Ascher SM, Arnold LL, Patt RH, Schrufer JJ, Bagley AS, Semelka RC, et al. Adenomyosis: Prospective comparison of MR imaging and transvaginal sonography. *Radiology* 1994;190:803-6.
 16. Reinhold C, Atri M, Mehio A, Zakarian R, Aldis AE, Bret PM. Diffuse uterine adenomyosis: Morphologic criteria and diagnostic accuracy of endovaginal sonography. *Radiology* 1995;197:609-14.
 17. Atzori E, Tronci C, Sionis L. Transvaginal ultrasound in the diagnosis of diffuse adenomyosis. *GynecolObstet Invest* 1996;42:39-41.
 18. Atri M, Reinhold C, Mehio AR, Chapman WB, Bret PM. Adenomyosis: US features with histologic correlation in an in-vitro study. *Radiology* 2000;215:783-90.
 19. Sloboda L, Molnar E, Popovic Z, Zivkovic S. Analysis of pathohistological results from the uterine mucosa 1965-1998 at the gynecology department in Senta. *Med Pregl* 1999;52:2635.
 20. S. Kayastha, Study of endometrial tissue in dysfunctional uterine bleeding , 1 March 2013, *Nepal Medical College journal : NMCJ* [10]
 21. S M Madan¹, Z A Al-Jufairi, 2001 Feb;22(2):153-6. Abnormal uterine bleeding. Diagnostic value of hysteroscopy, PMID: 11299412, *Saudi MedJ* [2]
 22. T Justin Clark¹, Doris Voit, Janesh K Gupta, Christopher Hyde, Fujian Song, Khalid S Khan, Accuracy of hysteroscopy in the diagnosis of endometrial cancer and hyperplasia: a systematic quantitative review, 2002 Oct 2; 288 (13) : 161021. doi: 10.1001/jama.288.13.1610. [3]
 23. Lithingo Lotha and Asha Borah, Sept 2016, *Clinicopathological evaluation of abnormal uterine bleeding in perimenopausal women* [7].
 24. Alev Ozer, Serdar Ozer, Mine Kanat-Pektas, Correlation between transvaginal ultrasound measured endometrial thickness and histopathological findings in Turkish women with abnormal uterine bleeding, 18 February 2016 [8]
 25. Qureshi FU, Yusuf AW. Distribution of causes of abnormal uterine bleeding using the new FIGO classification system. *JPMA*. 2013;63(8):973-75.