

A Comparative Evaluation of Palatal Rugae Pattern Among Male, Female and Transgender Population

¹Hantalacci D S, Dept. of Oral Pathology and Microbiology, School of Dental Sciences

²Bothova M V, Dept. of Oral Pathology and Microbiology, School of Dental Sciences

³Kambert F P, Dept. of Oral Pathology and Microbiology, School of Dental Sciences

Corresponding Author: Hantalacci D S, Dept. of Oral Pathology and Microbiology, School of Dental Sciences

Citation This Article: Hantalacci D S, Bothova M V, Kambert F P, “A Comparative Evaluation of Palatal Rugae Pattern Among Male, Female and Transgender Population”, IJHDC – January – February – 2025, Volume. – 4, Issue – 1, P. No. 04 – 09.

Open Access Article: This is an Open Access article that uses a funding model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.

Type of Publication: Original Research Article

Conflicts of Interest: Nil

Abstract

Introduction: The irregular ridges of the mucous membrane on the anterior portion of the palate are called palatal rugae, or palatine rugae. They differ in size, form, and arrangement, making each person unique. These rugae patterns begin during fetal development and last the entirety of an individual's life. Researchers are interested in palatal rugae patterns because of their clinical importance in domains such as forensic identification and orthodontics.

Aim: The purpose of this study was to compare the palatal rugae pattern among the male, female, and transgender populations.

Settings and Design: This study was cross-sectional and conducted on a convenience sample of 90 subjects selected. The study involved 30 males, 30 females, and 30 eunuchs in the age range of 17 to 35 years.

Materials and Methods: Maxillary impressions using alginate impression material were made, and the cast was prepared using a die stone on the palatal area and a dental stone as a base. The palatal rugae pattern was assessed based on number, length, shape, direction, and unification.

Statistical Analysis Used: One-way ANOVA was used to compare the mean values between different genders. The multiple pairwise comparisons were done with the Bonferroni post hoc correction. The statistical significance was fixed at 0.05.

Results: There were statistically significant differences in some parameters, like the number of rugae, fragmentary rugae, wavy rugae, curve rugae, forwardly directed rugae, and backwardly directed rugae, between transgender and other gender groups.

Conclusion: The difference in the parameters of the palatal rugae pattern between the transgender population

and the other gender group is attributed to genetic makeup and sexual dimorphism.

Keywords: Sexual Dimorphism, Fragmentary Rugae, Wavy Rugae, Curve Rugae, Forensic Identification and Orthodontics.

Introduction

Palatal rugae, also known as palatine rugae, are amorphous ridges of mucous membrane on the front region of the palate. Every individual is different due to their variations in shape, size, and arrangement. Rugae patterns appear during prenatal development and continue throughout an individual's life. Palatal rugae patterns are of interest to researchers due to their therapeutic value in fields like forensic identification and orthodontics. It has been noted that the sexes have different palatal rugae patterns. Studies have shown that the patterns of palatal rugae differ in men, women, and transgender individuals. Research has shown that there are statistically significant differences between males, women, and transgender individuals palatal rugae patterns. Disparities in the Patterns of Palatal Rugae by Gender. Based on available data, palatal rugae patterns seem to vary across men, females, and transgender individuals.

But it's important to keep in mind that there are other variables than heredity that contribute to the differences in palatal rugae patterns amongst males. Environmental factors, such as oral habits, may also lead to variations in palatal rugae patterns. Male palatal rugae patterns have been analyzed, and some distinctive characteristics have been discovered. These characteristics include variations in the rugae patterns' size, shape, and arrangement. Males tend to have bigger, more asymmetrical rugae patterns than females. Additionally, males often have a greater number of rugae patterns than females.

Consequently, while establishing someone's ethnicity based on palatal rugae patterns, it is crucial to take a variety of things into account.

Because of their special characteristics that last a lifetime, the palatal rugae patterns have been used as reference locations for superimposition both before and after orthodontic treatment. Similar to fingerprints, they are also useful tools for forensic identification due to their uniqueness. It is nearly always necessary to use forensic methods to ascertain the identity of transgender people during catastrophes since the forensic literature has focused mostly on the two genders of men and women.

Methodology

This cross-sectional research was done on a suitable sample of 90 people. The research included 30 men, 30 females, and 30 transgender people, ranging in age from 17 to 35 years. The selection of the research participants was done according to inclusion and exclusion standards.

Inclusion criteria

- The individuals' ages were chosen to be between 17 and 45 years old.
- Individuals without prosthesis.

Exclusion criteria

- Individuals with palate and lip anomalies, such as cleft palate and cleft lip
- Individuals using prostheses
- People receiving orthodontic treatment.

After obtaining informed permission, maxillary imprints were produced using alginate impression material, and research models were created using die stone and dental stone. There were no air bubbles or cavities in any of the chosen casts, particularly in the front third of the palate. By using a black marker pen to outline the rugae pattern

on the research models, the pattern's clarity on the cast was improved. The following heading applies to this categorization.

Considering the number of palatal rugae: The total number of rugae on the left and right sides of the mid-palatine raphe was counted. The rugae were then divided into three kinds according to their length, which are listed below: Primary >5 mm, Secondary 3 to 5 mm, and Fragmentary <3 mm. Less than 2 mm of rugae were ignored. The length of a rugae was established by measuring its biggest dimension, independent of form, using a divider. Using a stainless steel scale, the measurement on the divider was then converted to millimeters.

Rugae unification: When two rugae are linked at their origin or end, it is stated that a rugae is unified. Based on the unification, two groups were created for the rugae.

Diverging: When two rugae originate from the midline but branch off right away Rugae that diverge from the midline but converge at their lateral section are referred to as being convergent. One qualified and calibrated assessor assessed the palatal rugae pattern. To establish intra examiner reliability, the examination was done again on 10 randomly chosen casts separated by 3 hours.

Statistical analysis

Data was entered into the Excel sheet. Data were analyzed using SPSS (Statistical Package for Social Sciences) 25.0 version, IBM, Chicago. Descriptive statistics were performed, and data were described as mean ± standard deviation. The inter-group comparison was done using the One-way ANOVA followed by post hoc analysis, if needed. P-value<.05 was considered statistically significant.

Results

The study had 90 subjects, including 30 males, 30 females, and 30 transgenders.

Total rugae

The mean ± standard deviation of the total number of rugae among males, females, and transgender was 10.7667 ± 1.45468, 10.4667 ± 1.88887, and 9.3667 ± 1.35146 respectively. The difference in the mean of the total number of rugae was statistically significant (p-value<.05). Post hoc analysis revealed that the number of rugae was significantly less among transgenders compared to males and females (p-value <.05) whereas the difference in the number of rugae between males and females was statistically non-significant (p-value>.05).

Table 1: Comparison of the median of the total number of rugae between three groups

	Male	Female	Transgender	F-value	p-value
Mean ± standard deviation	10.7667 ± 1.45468	10.4667 ± 1.88887	9.3667 ± 1.35146	6.511	.002*

One-way ANOVA. *p-value<.05 was considered statistically significant.

Table 2: Post hoc analysis

Pair-wise	p-value
Male vs female	.744
Male vs transgender	.003*
Female vs transgender	.023*

*p-value<.05 was considered statistically significant.

Length of rugae

The mean ± standard deviation number of primary rugae was highest in males followed by females followed by transgenders. The difference in the number of primary and secondary rugae between the groups was statistically non-significant (p-value >.05). The number of fragmentary rugae was found to differ significantly

between the groups (p-value<.05). Post hoc analysis revealed that the number of fragmentary rugae was significantly less among transgenders compared to male (p-value <.05). However, the difference in the number of fragmentary rugae between males and females, and females and transgenders were statistically non-significant (p-value >.05).

Table 3: Comparison of the median length of rugae between three groups

		Male	Female	Transgender	F- value	p-value
Mean ± standard deviation	Primary	5.7333 ± 3.12866	5.3333 ± 2.75848	5.0333 ± 2.98829	.422	.657
	Secondary	2.8667 ± 2.94470	2.9000 ± 2.63072	3.1000 ± 2.59110	.064	.938
	Fragmentary	2.0667 ± 2.04995	2.2333 ± 2.17641	1.0333 ± 1.03335	3.801	.026*

One-way ANOVA. *p-value <.05 was considered statistically significant.

Table 4: Post hoc analysis

Pair-wise	P-value
Male vs female	.934
Male vs transgender	.078
Female vs transgender	.034*

*p-value<.05 was considered statistically significant.

Shape of rugae

The mean ± standard deviation of the number of straight, wavy, circular, divergent, and convergent rugae was not significantly different between the groups (p-value >.05). However, the number of curve rugae was significantly

different between the groups (p-value <.05). Post hoc analysis revealed that the number of curve rugae was significantly less among transgender compared to females (p-value <.05) however, the difference in the number of curve rugae was non-significant between males and females and, males and transgenders (p-value >.05).

Table 5: Comparison of shape of rugae among subjects belonging to three groups

		Male	Female	Transgender	F-value	p-value
Mean ± standard deviation	Straight	2.7000 ± 1.20773	2.8667 ± 1.13664	3.0667 ± 1.01483	.802	.452
	Wavy	2.5333 ± 1.35782	2.5333 ± 1.22428	2.0000 ± 1.50860	1.519	.225
	Circular	2.4000 ± 1.13259	1.8667 ± 1.13664	1.9667 ± 0.80872	2.240	.113
	Curve	1.4333 ± 1.22287	1.8000 ± 1.09545	1.0667 ± 0.86834	3.508	.034*
	Divergent	1.0333 ± 0.85029	0.9667 ± .63335	0.7000 ± 0.54971	1.190	3.09
	Convergent	0.5333 ± 0.33030	0.4333 ± 0.27891	0.5333 ± 0.38145	.206	.814

One-way ANOVA. *p-value <.05 was considered statistically significant.

Table 6: Post hoc analysis

Pair-wise	p-value
Male vs female	.386
Male vs transgender	.386
Female vs transgender	.026*

*p-value<.05 was considered statistically significant.

Discussion

According to the study's findings, palatal rugae patterns are specific to each person and can be a useful genetic research tool. Additionally, prior studies have demonstrated the existence of genetic connections in rugae patterns, which vary between genders and races.

Palatal Rugae Patterns among Males Several unique traits have been found when analyzing palatal rugae patterns among males. These features include differences in the size, form, and arrangement of the rugae patterns. In comparison to females, males often have larger, more asymmetrical rugae patterns. Furthermore, compared to females, males frequently display a higher number of rugae patterns.

The purpose of the study was to ascertain whether females' palatal rugae patterns differed from those of other genders in any notable way. The palatal rugae patterns of different females did differ, the researchers discovered. Compared to males and transgender people, females have different numbers, shapes, and lengths of palatal rugae. According to the study, males had a larger frequency of curved and wavy palatal rugae, while females had a greater prevalence of straight-shaped rugae. ⁸The number, length, and direction of the major palatal rugae did not significantly differ between the male and female groups, according to the study. The results of this study imply that

palatal rugae patterns could not be a good way to identify men, females, and transgender people in this community.

These results are in line with earlier studies that propose a link between gender differentiation and palatal rugae patterns.

To make definitive conclusions on the relationship between palatal rugae patterns and gender differentiation, the results of these studies emphasize the need for additional research and consideration of aspects including ethnic variance, sample size, and study population. Palatine rugae's resilience and uniqueness have made them useful for forensic identification. In contrast, males had more instances of wavy and curved rugae. It is noteworthy that a more thorough investigation of gender variations in palatal rugae form changes would be possible with a larger sample size analysis.

Additionally, when comparing male versus female subjects, differences were seen across secondary rugae types, such as fragmentary, straight, and perpendicular-directed. Our findings showed that female participants had significantly higher rugae counts.

Conclusion

The difference in the mean of the total number of rugae was statistically significant (p-value<.05).Post hoc analysis revealed that the number of rugae was significantly less among transgenders compared to males and females (p-value <.05).The number of fragmentary

rugae was found to differ significantly between the groups (p -value $<.05$). Post hoc analysis revealed that the number of fragmentary rugae was significantly less among transgender compared to male (p -value $<.05$). Post hoc analysis revealed that the number of curve rugae was significantly less among transgender compared to females (p -value $<.05$)

In summary, the identification of palatal rugae patterns among different genders has wide-ranging implications for public health, including improved identification methods, more effective healthcare services, and enhanced inclusivity. Transgender people can be identified based on self-identification. These types of studies can help to establish the identification between transgender and the individuals who are imposing to be transgender. These considerations contribute to a more comprehensive and equitable public health approach that serves the diverse needs of the population.

References

1. Bansode SC, Kulkarni MM. Importance of palatal rugae in individual identification. *J Forensic Dent Sci.* 2009;1:77–81.
2. Mujoo S, Sakarde S, Sur J, Singh A, Khan F, Jain S, et al. Cheiloscropy and Paltoscropy: Anovel tool for sex identification. *Chettinad Health City Med J.* [Last accessed on 2012 Aug 29]. pp. 146–50.
3. Filho Ismar EM. Palatal rugae patterns as bioindicators of identification in Forensic Dentistry. *RFO.* 2009;14:227–33.
4. Eboh DE. Palatal Rugae Patterns of Urhobos in Abraka, South-Southern Nigeria. *Int J Morphol.* 2012;30:709–13.
5. Lysell L. Plicae palatinae transversae and papilla incisiva in man. A morphologic and genetic study. *Acta Odontol Scand.* 1955;13:5–137.
6. Venegas VH, Valenzuela JS, Lopez MC, Galdames IC. Palatal rugae: systematic analysis of its shape and dimensions for use in human identification. *Int J Morphol* 2009 Sep;27(3):819-825.
7. Muhasilovic S, Hadziabdic N, Galic I, Vodanovic M. Analysis of palatal rugae in males and females of an average age of 35 in a population from Bosnia and Herzegovina (Sarajevo Canton). *J Forensic Leg Med* 2016 Apr;39:147-150.
8. Silva-Sousa AC, Marañón-Vásquez GA, Stuaní MBS, Proff P, Andrades KMR, Baratto-Filho F, Matsumoto MAN, Paddenberg E, Kúchler EC, Kirschneck C. Genetic variants in bone morphogenetic proteins signaling pathway might be involved in palatal rugae phenotype in humans. *Scientific Reports.* 2021;11(1): 12715.
9. Tey SN, Lin YM, Mohamed AMFS. Stability of palatal rugae after orthodontic/orthopaedic expansion: A scoping review. *Australasian Orthodontic Journal.* 2023;39(1):158–170.
10. Moran A, Tippett H, Manoharan A, Cobourne MT. Alteration of palatine ruga pattern in subjects with oligodontia: A pilot study. *American Journal of Orthodontics and Dentofacial Orthopedics,* 2016;150(2), 295–302.
11. Trizzino A, Messina P, Sciarra FM, Zerbo S, Argo A, Scardina GA. Palatal Rugae as a Discriminating Factor in Determining Sex: A New Method Applicable in Forensic Odontology? *Dentistry Journal.* 2023;11 (9):204.
12. Gezer R, Deniz M, Uslu AI. Morphological characteristics and individual differences of palatal rugae. *Journal of Craniofacial Surgery.* 2019;30(6):1906–1910.